

BtsC I

Cat. No.	용량	농도
DYR1270	2,000 units	20 units/μl
DYR1272	4,000 units	20 units/μl
DYR1274	10,000 units	20 units/μl
DYR1276	10,000 units	100 units/μl

◆ **제품구성**

- BtsC I
- 10X DY Buffer IV
- 10X FastCut Buffer
- Sterile water
- Dyne 6X DNA Loading Buffer ver.2

◆ **Source**

- *Bacillus thermosphaericus*

◆ **Quality control**

- Unit definition assay
- Overdigestion assay
- Endonuclease assay
- Extreme purity assay

◆ **인식부위**



Single letter code

W = A or T	S = C or V = A or C or G
N = A or C or G or TG	M = A or C
K = G or T	R = A or G
Y = C or T	B = C or G or T
D = A or G or T	H = A or C or T

◆ **보관온도**

- -20°C

◆ **Heat inactivation**

- 80°C for 20 min.

◆ **Unit정의**

- 1 unit은 박테리오파지 λ DNA 1 μg을 50 μl 반응물로 60°C에서 1시간 동안 완전히 분해하는데 필요한 효소의 양이다.

◆ **Buffer별 상대적 활성도**

I	II	III	IV	FastCut
75%	100%	100%	100%	100%

◆ **Methylation effect**

Methylation	<i>dam</i>	<i>dcm</i>	CpG
Cleavage	Cleavage	Cleavage	Cleavage

◆ **주의사항**

- BtsC I은 BstF5 I의 동일서열인식 제한효소이다. *dam*, *dcm* 또는 CpG 메틸화(methylation)의 영향을 받지 않는다.

◆ **표준반응 조건**

- Normal Protocol

Component	농도	Volume
Substrate DNA	1 μg	X μl
10X DY Buffer IV	1 X	5 μl
BtsC I		Substrate dependent
Sterile water		Up to 50 μl

- * Incubate at 50°C for 1 hr

- Fast Protocol

Component	농도	Volume
Substrate DNA	1 μg	X μl
10X FastCut Buffer	1 X	5 μl
BtsC I	20 unit	1 μl
Sterile water		Up to 50 μl

- * Incubate at 50°C for 15 min